



Future Circular Colliders project (FCC): a long term vision for Particle Physics

Stéphane Monteil,
Clermont University, LPC-IN2P3-CNRS.

Outline

- The SM in a glance and the open questions.
- Introduction to the Future Circular Colliders project.
- The electron machine and some prototype detectors.
- Physics case at large.
- Implementation as an outlook.

The free parameters of the SM:

- $SU(2)_L \otimes U(1)_Y$ unification:
 - the weak and electromagnetic coupling constants G_F/g_W and α_{EM} .
- After the spontaneous breaking of the symmetry:
 - The nine masses of the fermions: m_f .
 - The masses of the electroweak gauge bosons: m_Z and m_W .
 - The scalar sector parameters: $V(\phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$
 v (the v.e.v) and m_H .

1) Scientific context: SM became a theory



The free parameters of the SM

- The CKM matrix elements : it's a 3X3 complex and unitary matrix and hence can be described by means of only **4 independent parameters**. As the masses of the fermions (except for the top quark), these 4 parameters are decoupled from the rest of the theory.
- If you like QCD in (and you do), just add α_S (and θ^S_{CP}).
- Neutrino oscillations are implying neutrinos to be massive and to mix \rightarrow 7 parameters to minimally describe them.
- The number of parameters amounts to 20 (28 w/ neutrinos and strong CP). Not all of them are independent though.

1) Scientific context: SM became a theory

Reorganisation:

- QCD and α_s : LEP and others did great already. Limitation of the consistency test is not yet fully on the theory side for most of the determinations.




- A better α_s determination is desirable, and in order for advanced predictions (QCD x-sections, Higgs decays, top mass, Z width, R_b , R_l).

1) Scientific context: SM became a theory

Reorganisation:


- The nine masses of the fermions: m_f .
- They are for 8 of them decoupled from the rest of the SM parameters.
- Nothing much to do here as well till the moment a theory comes with a prediction.
- The top quark has a specific status because it enters dominantly in the radiative corrections of the intermediate bosons mass propagators (in particular), *e.g.*



1) Scientific context: SM became a theory

Reorganisation:




- The (4) CKM matrix elements (decoupled from the rest of the theory). The consistency check of the SM hypothesis in that sector is a pillar of the SM:



1) Scientific context: SM became a theory

Reorganisation:



- The rest of the free parameters are part of the so-called electroweak precision observables consistency check. This is the other pillar of the SM. Fix G_F , α_{EM} and m_Z at their measured value and produce a prediction of m_{top} , m_W and m_H . A tremendous success !



1) Scientific context: SM became an (invincible) theory

Recap:


- Two pillars: EWPT and Flavours.



1) Scientific context: SM became an (invincible) theory

Recap:



- Two pillars: EWPT and Flavours.



1) Scientific context: SM became an (invincible) theory

Recap:



- Two pillars: EWPT and Flavours.



1) Scientific context: SM became an (invincible) theory

Recap:

- Two pillars: EWPT and Flavours



1) Scientific context: SM became an invincible theory

Lessons

- The SM has cleared so far the attacks from LEP, TeVatron, *B*-factories, LHC and single-observables experiments.
- There are compelling beauty arguments for Beyond Standard Model (BSM) Physics. I will overlook them.
- Instead, three indisputable measurements/observations are crying for BSM:
 - The neutrinos have a mass. Though several ways exist theoretically, it's tempting / natural to enhance the neutral particle content with right-handed states.
 - Dark matter: a nice (recent-ish) evidence for cosmological dark matter is the observation of a low surface brightness galaxy [ArXiv:1606.06291].
 - Baryonic asymmetry in the Universe.

1) Scientific context: scenarii

1) Find a new heavy particle at the Run III of LHC:

- HL-LHC can study it to a certain extent.
- If mass is small enough (and couples to electrons), CLIC can be the way.
- Larger energies are needed to study (find) the whole spectrum.
- The underlying quantum structure must be studied.

2) Find no new particle, but non-standard H properties

- HL-LHC can study it to a certain extent.
- Higgs factory.
- Z , W , top factories for the quantum structure.
- Energy frontier (also for precision measurements)

3) Find no new particle, standard H properties but flavour observables departing from SM:

- Z , W , top factories for the quantum and flavour structure.
- Energy frontier to find the corresponding spectrum.

4) Find no new particle, standard H properties and flavour observables in SM:

- Asymptotic Z , W , H , top factories for asymptotic precision.
- Push the energy frontier to the best of our knowledge.

1) Scientific context: scenarii

1) Find a new heavy particle at the Run III of LHC:

- HL-LHC can study it to a certain extent.
- If mass is small enough (and couples to electrons), CLIC can be the way.
- Larger energies are needed to study (find) the whole spectrum.
- The underlying quantum structure must be studied.

2) Find no new particle, but non-standard H properties

- HL-LHC can study it to a certain extent.
- Higgs factory.
- Z , W , top factories for the quantum structure.
- Energy frontier (also for precision measurements)

3) Find no new particle, standard H properties but flavour observables departing from SM:

- Z , W , top factories for the quantum and flavour structure.
- Energy frontier to find the corresponding spectrum.

4) Find no new particle, standard H properties and flavour observables in SM:

- Asymptotic Z , W , H , top factories for asymptotic precision.
- Push the energy frontier to the best of our knowledge.

1) Scientific context: scenarii by anticipation of what follows

1) Find a new heavy particle at the Run III of LHC:

- HL-LHC can study it to a certain extent.
- If mass is small enough (and couples to electrons), CLIC can be the way.
- Larger energies are needed to study (find) the whole spectrum [FCC-hh].
- The underlying quantum structure must be studied [FCC-ee].

2) Find no new particle, but non-standard H properties

- HL-LHC can study it to a certain extent.
- Higgs factory [ILC, FCC-ee].
- Z , W , top factories for the quantum structure [FCC-ee].
- Energy frontier (also for precision measurements) [FCC-hh].

3) Find no new particle, standard H properties but flavour observables departing from SM:

- Asymptotic Z , W , top factories to fix the energy scale [FCC-ee].
- Energy frontier to find the corresponding spectrum [FCC-hh].

4) Find no new particle, standard H properties and flavour observables in SM:

- Asymptotic Z , W , H , top factories for asymptotic precision [FCC-ee].
- Push the energy frontier to the best of our knowledge [FCC-hh].

1) Scientific context: scenarii by anticipation of what follows

1) Find a new heavy particle at the Run III of LHC:

- HL-LHC can study it to a certain extent.
- If mass is small enough (and couples to electrons), CLIC can be the way.
- Larger energies are needed to study (find) the whole spectrum [FCC-hh].
- The underlying quantum structure must be studied [FCC-ee].

2) Find no new particle, but non-standard H properties

- HL-LHC can study it to a certain extent.
- Higgs factory [ILC, FCC-ee].
- Z , W , top factories for the quantum structure [FCC-ee].
- Energy frontier (also for precision measurements) [FCC-hh].

3) Find no new particle, standard H properties but flavour observables departing from SM:

- Asymptotic Z , W , top factories to fix the energy scale [FCC-ee].
- Energy frontier to find the corresponding spectrum [FCC-hh].

4) Find no new particle, standard H properties and flavour observables in SM:

- Asymptotic Z , W , H , top factories for asymptotic precision [FCC-ee].
- Push the energy frontier to the best of our knowledge [FCC-hh].

1) From Scientific context to the introduction to FCC

2. Introduction to FCC project:

- Starting from the former European HEP strategy 2013

Summary: European Strategy Update 2013 *Design studies and R&D at the energy frontier*

....“to propose an ambitious **post-LHC accelerator project at CERN** by the time of the next Strategy update”:

- d) **CERN should undertake design studies for accelerator projects in a global context,**
 - *with emphasis on **proton-proton and electron-positron high-energy frontier machines.***
 - *These design studies should be coupled to a vigorous accelerator **R&D programme, including high-field magnets and high-gradient accelerating structures,***
 - **in collaboration with national institutes, laboratories and universities worldwide.**
 - <http://cds.cern.ch/record/1567258/files/esc-e-106.pdf>




- At the time the LHC Run II will have delivered a significant part of its results, have an educated vision of the reach of future machines for the next round of **the European Strategy in 2020.**

2. Introduction to FCC: the scope of the project


Forming an international coll.
(hosted by Cern) to study:

- 100 TeV pp -collider (FCC- hh) as long term goal, defining infrastructure requirements.
- e^+e^- collider (FCC- ee) as potential first step.
- $p-e$ (FCC- he) as an option.
- 80-100 km infrastructure in Geneva area.
- Conceptual design report and cost review for the next european strategy → 2020.




2. Introduction to FCC: the completion of the CDR


The Design Study is completed and fulfilled the mandate



©M. Benedikt



Results of FCC Conceptual Design Study



more than 1350 contributors from 350 institutes, a truly global collaboration and effort as suggested by the EPPSU 2013

Study Documentation:

4 CDR volumes submitted to EPJ in December 2018.

- FCC Physics Opportunities
- FCC-ee
- FCC-hh
- HE-LHC

Preprints available since 15 January 2019
<http://fcc-cdr.web.cern.ch/>

CDR presentation during welcome event this evening.

Paper copies can be requested at
<http://get-fcc-cdr.web.cern.ch>

Next stages (CDR+, TDR) are subjected to the ESPP update outcome.

2. Introduction to FCC: the ESPP update

Three recommendations:

- *The particle physics community should ramp up its R&D effort focused on advanced accelerator technologies, in particular that for high-field superconducting magnets, including high-temperature superconductors;*
 - ➔ **To realize a machine at the energy frontier, high field magnets with at least 16T are mandatory and far from industrialisation, development of HTS magnets reaching higher fields should be pursued**
- *Europe, together with its international partners, should investigate the technical and financial feasibility of a future hadron collider at CERN with a centre-of-mass energy of at least 100 TeV and with an electron-positron Higgs and electroweak factory as a possible first stage. Such a feasibility study of the colliders and related infrastructure should be established as a global endeavour and be completed on the timescale of the next Strategy update.*
 - ➔ **Feasibility study should be carried out before the next Strategy Update to allow for decision to be taken**
 - **technical feasibility, administrative implications and questions of implementation in the Geneva area including tunnelling and environmental impact)**
 - **financial feasibility for construction and operation, including additional resources from international partners and start establishing the global frame for the project.**
- *The timely realisation of the electron-positron International Linear Collider (ILC) in Japan would be compatible with this strategy and, in that case, the European particle physics community would wish to collaborate.*

© U. Bassler

Note that everything is written in such a way that proponents of each and every large scale projects can read it happily. Yet,

“the vision is to prepare a Higgs factory, followed by a future hadron collider with sensitivity to energy scales an order of magnitude higher than those of the LHC ” ESPP 2020 d’après d’Hondt.



2. Introduction to FCC: same scope after ESPP



The FCC integrated program inspired by successful LEP – LHC programs at CERN

Comprehensive cost-effective program maximizing physics opportunities

- **Stage 1: FCC-ee (Z, W, H, $t\bar{t}$) as Higgs factory, electroweak & top factory at highest luminosities**
- **Stage 2: FCC-hh (~100 TeV) as natural continuation at energy frontier, with ion and eh options**
- Complementary physics
- Common civil engineering and technical infrastructures
- Building on and reusing CERN's existing infrastructure
- FCC integrated project allows seamless continuation of HEP after HL-LHC



2. Additional comments:


“the vision is to prepare a Higgs factory, followed by a future hadron collider with sensitivity to energy scales an order of magnitude higher than those of the LHC ” ESPP 2020 d’après d’Hondt.

- FCC-*ee* can’t happen before the completion of the HL-LHC program.
- FCC-*ee* shall happen seamlessly after HL-LHC.
- FCC-*hh* is foreseen after FCC-*ee*. The Higgs Physics program is inclusive (FCC-*hh* invincible for trilinear H couplings). The Flavour Physics program at FCC-*hh* is simultaneously appealing but subjected to in depth studies in light of LHCb U2.
- Link to the slide.
- Focus on the electron machine in the following.

2) From Introduction to FCC to the machine elements & detectors

3. The FCC e^+e^- machine. Baseline design


- Physics from the Z pole to top pair production (90 - 400 GeV), crossing WW and ZH thresholds with unprecedented statistics everywhere.
- Two rings (top-up injection) to cope with high current and large number of bunches at operating points up to ZH .
- Description of the machine parameters (given next slide).
- To some extent, SuperKEKB is already meeting or about to meet some of the challenges of FCC-ee:



Some SuperKEKB parameters :	
β_y^* : 300 μm	FCC-ee (H) : 1 mm
σ_y : 50 nm	FCC-ee (H) : 50 nm
ϵ_y/ϵ_x : 0.25%	FCC-ee (H) : 0.2% to 0.1%
e^+ production rate : $2.5 \times 10^{12} / \text{s}$	FCC-ee (H) : $< 1 \times 10^{11} / \text{s}$
Off-momentum acceptance at IP : $\pm 1.5\%$	FCC-ee (H) : $\pm 2.0\%$ to $\pm 2.5\%$
Beam Lifetime : 5 minutes	FCC-ee (H) : 20 minutes
Centre-of-mass energy: ~ 10 GeV	FCC-ee (H) : 240 GeV


3. The FCC e^+e^- machine. Baseline design



 FCC-ee collider parameters (stage 1)				
parameter	Z	WW	H (ZH)	ttbar
beam energy [GeV]	45	80	120	182.5
beam current [mA]	1390	147	29	5.4
no. bunches/beam	16640	2000	393	48
bunch intensity [10^{11}]	1.7	1.5	1.5	2.3
SR energy loss / turn [GeV]	0.036	0.34	1.72	9.21
total RF voltage [GV]	0.1	0.44	2.0	10.9
long. damping time [turns]	1281	235	70	20
horizontal beta* [m]	0.15	0.2	0.3	1
vertical beta* [mm]	0.8	1	1	1.6
horiz. geometric emittance [nm]	0.27	0.28	0.63	1.46
vert. geom. emittance [pm]	1.0	1.7	1.3	2.9
bunch length with SR / BS [mm]	3.5 / 12.1	3.0 / 6.0	3.3 / 5.3	2.0 / 2.5
luminosity per IP [$10^{34} \text{ cm}^{-2}\text{s}^{-1}$]	230	28	8.5	1.55
beam lifetime rad Bhabha / BS [min]	68 / >200	49 / >1000	38 / 18	40 / 18

- There seem to be several machines actually.

3. The FCC e^+e^- machine. Luminosity figure



- The FCC-ee offers the largest luminosities in its whole energy range.
- We're speaking here of 10^5 Z/s , 10^4 W/h, $1.5 \cdot 10^3$ H and top /d, in a very clean environment: no pile-up, controlled beam backgrounds, E and p constraints, without trigger. In particular, **you do the LEP in a minute!**

3. The FCC e^+e^- machine. Time allocation at Z pole

- The time / energy allocation of the machine has been worked out ...



Working point	Lumi. / IP [$10^{34} \text{ cm}^{-2} \cdot \text{s}^{-1}$]	Total lumi. (2 IPs)	Run time	Physics goal
Z first phase	100	26 ab^{-1} /year	2	
Z second phase	200	52 ab^{-1} /year	2	150 ab^{-1}

- ... we're speaking here of $5 \cdot 10^{12}$ Z, 10^8 WW, 10^6 H and 10^6 top pairs.
- Of particular relevance for the Flavour Physics is the Z pole (2 IPs — 4 are considered):

Particle specie at FCC- ee	B^0	B^+	B_s^0	Λ_b	B_c^+	$c\bar{c}$	$\tau^-\tau^+$
Yield ($\times 10^9$) [for $5 \cdot 10^{12}$ Z]	310	310	75	65	1.5 [†]	600	180

- 20 times Belle II for B^0 and B^+
- Direct comparison with LHCb yields requires a mode-by-mode approach to take into account trigger and reconstruction efficiencies.

3. A word on FCC e^+e^- detectors




- Two designs have been studied so far (and used to assess performances)
- Robust towards performance, intricate MDI, beam backgrounds.
- The key point for all the Physics program is the **lightness** ...
- Personal note: FCC project aims at providing four detector proposals by 2026. Among those proposals, there is room for a dedicated design for Flavours, in particular for hadron identification, vertexing and calorimeter.

3) From machine & detectors to the Physics case / reach


Be it only for the accurate study of the Higgs-boson decays, an electron collider is the way to go.

© P. Janot




4) Physics case — Higgs chapter

- Two energy points (240 and 360 GeV) for the program




- Invincible precision on the absolute couplings and width. Interplay with HL-LHC.



Collider	HL-LHC	FCC-ee		
Luminosity (ab^{-1})	3	5 @ 240GeV	+1.5 @ 365GeV	+HL-LHC
Years	25	3	+4	-
$\delta\Gamma_H/\Gamma_H$ (%)	SM	2.7	1.3	1.1
$\delta g_{HZZ}/g_{HZZ}$ (%)	1.3	0.2	0.17	0.16
$\delta g_{HWW}/g_{HWW}$ (%)	1.4	1.3	0.43	0.40
$\delta g_{Hbb}/g_{Hbb}$ (%)	2.9	1.3	0.61	0.55
$\delta g_{Hcc}/g_{Hcc}$ (%)	SM	1.7	1.21	1.18
$\delta g_{Hgg}/g_{Hgg}$ (%)	1.8	1.6	1.01	0.83
$\delta g_{H\tau\tau}/g_{H\tau\tau}$ (%)	1.7	1.4	0.74	0.64
$\delta g_{H\mu\mu}/g_{H\mu\mu}$ (%)	4.4	10.1	9.0	3.9
$\delta g_{H\gamma\gamma}/g_{H\gamma\gamma}$ (%)	1.6	4.8	3.9	1.1
$\delta g_{Htt}/g_{Htt}$ (%)	2.5	-	-	2.4
BR_{EXO} (%)	SM (0.0)	<1.2	<1.0	<1.0

4) Physics case – Higgs chapter – Even the electron Yukawa?




- The name of the game is to control the beam energy to touch the natural width of the Higgs boson. Monochromatisation technique.
- Tells you how exquisite can be the luminosity.

4) Physics case – EWPT chapter (including WW and top thr.)

Table 3.1: Measurement of selected electroweak quantities at the FCC-ee, compared with the present precisions.

Observable	present value \pm error	FCC-ee Stat.	FCC-ee Syst.	Comment and dominant exp. error
m_Z (keV/c ²)	91186700 \pm 2200	5	100	From Z line shape scan Beam energy calibration
Γ_Z (keV)	2495200 \pm 2300	8	100	From Z line shape scan Beam energy calibration
R_Z^l ($\times 10^3$)	20767 \pm 25	0.06	0.2-1	ratio of hadrons to leptons acceptance for leptons
$\alpha_s(m_Z)$ ($\times 10^4$)	1196 \pm 30	0.1	0.4-1.6	from R_Z^l above [29]
R_b ($\times 10^6$)	216290 \pm 660	0.3	<60	ratio of bb to hadrons stat. extrapol. from SLD [30]
σ_{had}^0 ($\times 10^3$) (nb)	41541 \pm 37	0.1	4	peak hadronic cross-section luminosity measurement
N_ν ($\times 10^3$)	2991 \pm 7	0.005	1	Z peak cross sections Luminosity measurement
$\sin^2 \theta_W^{eff}$ ($\times 10^6$)	231480 \pm 160	3	2 - 5	from A_{FB}^{ll} at Z peak Beam energy calibration
$1/\alpha_{QED}(m_Z)$ ($\times 10^9$)	128952 \pm 14	4	small	from A_{FB}^{ll} off peak [20]
$A_{FB}^{b,0}$ ($\times 10^4$)	992 \pm 16	0.02	1-3	b-quark asymmetry at Z pole from jet charge
$A_{FB}^{pol,\tau}$ ($\times 10^4$)	1498 \pm 49	0.15	<2	τ polarisation and charge asymmetry τ decay physics
m_W (keV/c ²)	80350000 \pm 15000	600	300	From WW threshold scan Beam energy calibration
Γ_W (keV)	2085000 \pm 42000	1500	300	From WW threshold scan Beam energy calibration
$\alpha_s(m_W)$ ($\times 10^4$)	1170 \pm 420	3	small	from R_e^W [31]
N_ν ($\times 10^3$)	2920 \pm 50	0.8	small	ratio of invis. to leptonic in radiative Z returns
m_{top} (MeV/c ²)	172740 \pm 500	20	small	From tt threshold scan QCD errors dominate
Γ_{top} (MeV/c ²)	1410 \pm 190	40	small	From tt threshold scan QCD errors dominate
$\lambda_{top}/\lambda_{top}^{SM}$	1.2 \pm 0.3	0.08	small	From tt threshold scan QCD errors dominate
ttZ couplings	\pm 30%	<2%	small	From $E_{CM} = 365$ GeV run

Z pole
tt thr. WW thr.



- Ultimate quantum completeness consistency test of the SM.
- The improvements in theory prediction precision is part of the FCC program.


4) Physics case – EWPT chapter

Table 3.1: Measurement of selected electroweak quantities at the FCC-ee, compared with the present precisions.

Observable	present value \pm error	FCC-ee Stat.	FCC-ee Syst.	Comment and dominant exp. error
m_Z (keV/c ²)	91186700 \pm 2200	5	100	From Z line shape scan Beam energy calibration
Γ_Z (keV)	2495200 \pm 2300	8	100	From Z line shape scan Beam energy calibration
R_Z^f ($\times 10^3$)	20767 \pm 25	0.06	0.2-1	ratio of hadrons to leptons acceptance for leptons
$\alpha_s(m_Z)$ ($\times 10^4$)	1196 \pm 30	0.1	0.4-1.6	from R_Z^f above [29]
R_b ($\times 10^6$)	216290 \pm 660	0.3	<60	ratio of bb to hadrons stat. extrapol. from SLD [30]
σ_{had}^0 ($\times 10^3$) (nb)	41541 \pm 37	0.1	4	peak hadronic cross-section luminosity measurement
N_ν ($\times 10^3$)	2991 \pm 7	0.005	1	Z peak cross sections Luminosity measurement
$\sin^2 \theta_{eff}^e$ ($\times 10^6$)	231480 \pm 160	3	2 - 5	from A_{FB}^{ll} at Z peak Beam energy calibration
$1/\alpha_{QED}(m_Z)$ ($\times 10^9$)	128952 \pm 14	4	small	from A_{FB}^{ll} off peak [20]
$A_{FB}^{b,0}$ ($\times 10^4$)	992 \pm 16	0.02	1-3	b-quark asymmetry at Z pole from jet charge
$A_{FB}^{pol,\tau}$ ($\times 10^4$)	1498 \pm 49	0.15	<2	τ polarisation and charge asymmetry τ decay physics
m_W (keV/c ²)	80350000 \pm 15000	600	300	From WW threshold scan Beam energy calibration
Γ_W (keV)	2085000 \pm 42000	1500	300	From WW threshold scan Beam energy calibration
$\alpha_s(m_W)$ ($\times 10^4$)	1170 \pm 420	3	small	from R_ℓ^W [31]
N_ν ($\times 10^3$)	2920 \pm 50	0.8	small	ratio of invis. to leptonic in radiative Z returns
m_{top} (MeV/c ²)	172740 \pm 500	20	small	From tt threshold scan QCD errors dominate
Γ_{top} (MeV/c ²)	1410 \pm 190	40	small	From tt threshold scan QCD errors dominate
$\lambda_{top}/\lambda_{top}^{SM}$	1.2 \pm 0.3	0.08	small	From tt threshold scan QCD errors dominate
ttZ couplings	\pm 30%	<2%	small	From $E_{CM} = 365$ GeV run

Z pole

 tt thr. WW thr.



- Ultimate quantum completeness consistency test of the SM.
- The improvements in theory prediction precision is part of the FCC program. Precision 1.4 GeV.

4) Physics case — Miscellanea chapter; some favourites.

- Higgs, top and EWK precision physics do not saturate the case.
- Many BSM direct searches thanks to the exquisite luminosity.
- Here are direct searches for heavy neutral leptons.

Three Generations of Matter (Fermions) spin 1/2

	I	II	III	
mass	2.4 MeV	1.27 GeV	173.2 GeV	0
charge	2/3	2/3	2/3	0
name	u up	c charm	t top	g gluon
	Left	Left	Right	0
	Right	Right	Left	0
Quarks	4.8 MeV	104 MeV	4.2 GeV	0
	-1/3	-1/3	-1/3	0
	d down	s strange	b bottom	γ photon
	Left	Left	Right	0
	Right	Right	Left	0
	0	0	0	91.2 GeV
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	0
	N_1	N_2	N_3	Z weak force
	Left	Left	Right	0
	Right	Right	Left	0
Leptons	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	0
	e electron	μ muon	τ tau	W [±] weak force
	Left	Left	Right	0
	Right	Right	Left	0






FIG. 2. Typical decays of a neutral heavy lepton via (a) charged current and (b) neutral current. Here the lepton l_i denotes $e, \mu, \text{ or } \tau$.



updated plot from MaD et al [1609.09069](#)
and Antusch et al [1710.03744](#)
cf. also Cai et al [1711.02180](#)

4) Physics case — Miscellanea chapter; some favourites.

- Higgs, top and EWK precision physics do not saturate the case.
 - Many BSM direct searches thanks to the exquisite luminosity.
 - Lepton Flavour violating Z decays.
-
- Lepton Flavour-Violating Z decays **in the SM** with lepton mixing are typically $< 10^{-50}$.
 - **Any observation** of such a decay would be an **indisputable evidence for New Physics**. FCC- ee exploration [JHEP 1504 (2015) 051]. $Z \rightarrow \tau\mu/e$ is unique at FCC.
 - The dominant background is ($Z \rightarrow \tau\tau$), where one tau decays into a close to beam energy lepton. The search is limited by the momentum resolution. A lot of phenomenology to explore yet.




Bottomline: With the expected tracking performance at FCC- ee (beam spread equivalent resolution at 45 GeV), the current limits are pushed by three orders of magnitude.

4) Physics case — Miscellanea chapter; some favourites.

- Flavour physics is a must do (remember Marie-Hélène's discussion) — FCC as ultimate Flavour factory the hierarchy does not tell the importance:
 - 1) Rare b -flavoured particles decays (EWP & friends).
 - 2) Di-leptonic decays (*e.g.* $B^0 \rightarrow \mu^+\mu^-$, $B_s \rightarrow \tau^+\tau^-$).
 - 3) (Semi-)leptonic decays (*e.g.* R_{D,D^*} , to $B_c \rightarrow \tau^+\nu \dots$)
 - 4) **CP violation study program at large.**
 - 5) Mass and lifetime properties, spectroscopy.
 - 6) Charm physics. Heavy Flavours Production

4) Physics case — Miscellanea chapter; some favourites.

A flavour of the potential: note the exquisite invariant-mass resolution. The probability of a pion to punch through and be identified as muon is taken as ALEPH performance.



© FCC EPJ+ essay

4) Heavy Flavours: physics opportunities conclusions






- FCC-ee gathers most of the advantages of each of the Flavour factories environment (pp, Upsilon).
- In turn, lots of physics opportunities and some likely unique physics cases and reach.
- They can't be met with any detector performance (vertexing beyond state-of-the-art, high resolution ECAL, PID).
- In particular if 4 IP are to be designed, it can be conceived that one experiment has a flavour-oriented design.
- Assessing the detector performance is the name of the game for the next five years.
- Convenient simulation tools have been set up. The fun can start.

5. The FCC implementation as an outlook





5. The FCC implementation: civil engineering.

- Machine footprints, experimental caverns, geological studies




5. The FCC implementation: timeline.

- Eighteen years towards Physics. No overlap in Physics between the end of HL-LHC and FCC-ee



5. The FCC implementation: timeline.

- Eighteen years towards Physics. No overlap in Physics between the end of HL-LHC and FCC-ee. The big picture.



- Is it crazy to plan a Physics program for seventy years?

5. The FCC implementation: timeline

- Is it reasonable to plan a Physics program for seventy years? It was.
- The previous HEP European planning was only for ... 60 years !

PHYSICS WITH VERY HIGH ENERGY

e^+e^- COLLIDING BEAMS

CERN 76-18
8 November 1976

L. Camilleri, D. Cundy, P. Darriulat, J. Ellis, J. Field,
H. Fischer, E. Gabathuler, M.K. Gaillard, H. Hoffmann,
K. Johnsen, E. Keil, F. Palmonari, G. Preparata, B. Richter,
C. Rubbia, J. Steinberger, B. Wiik, W. Willis and K. Winter

ABSTRACT

This report consists of a collection of documents produced by a Study Group on Large Electron-Positron Storage Rings (LEP). The reactions of


5. The FCC implementation — Cost



FCC-ee cost estimate


Total construction cost phase1 (Z, W, H) amounts to 10,500 MCHF

- 5,400 MCHF for civil engineering (51%)
- 2,000 MCHF for technical infrastructure (19%)
- 3,100 MCHF accelerator and injector (20%)



Complement cost for phase2 (tt) amounts to 1,100 MCHF

- 900 MCHF for RF, 200 MCHF for associated technical infrastructure




5. The FCC implementation — Future (future)



FCC-hh cost estimate


Total construction cost in “stand-alone” is 24,000 MCHF

- 13,600 MCHF accelerator and injector (57%)
 - Major part corresponds to the 4,700 Nb₃Sn 16 T main dipole magnets, totalling 9,400 MCHF, at cost target of 2 MCHF/magnet.
- 6,000 MCHF construction cost for surface and underground civil engineering (25%)
- 4,400 MCHF for technical infrastructures (18%)




Total construction cost in “combined mode” following FCC-ee is 17,000 MCHF.

- CE and TI from FCC-ee re-used
- 600 MCHF for additional CE structures:
 - Two experiment caverns for the lower luminosity experiments
 - Beam dump tunnels and the two transfer lines from LHC
- 2,800 MCHF for additional TI, driven by cryogenics infrastructure



6) Summary

- Achille Stocchi told you about the brilliant ERL technology to come. Here is what American colleagues thought of for FCC-ee. Check the top threshold. HHH is at hand ! Or we get modest at power consumption !



6) Summary

- The project is mature. FCC can be done ! The FCC software and detector full simulations are getting up. Please check:
 - <https://hep-fcc.github.io/FCCeePhysicsPerformance/>
- Fantastic tool for Higgs, top, and EWPT tests.
- The Flavour Physics case is now part of the FCC program in its own right. There are exciting times ahead with Flavours with the present data and this to come. It can be a lot of fun to think of these perspectives and some serious studies to be conducted. FCC-*ee* precision shall meet or increase the precision of each and both of Belle II and LHCb upgrades (super-complementarity).
- There are opportunities for internships at FCC-*ee* during the feasibility phase.

00) Back-up slides

Wish list for today:


- FCC tunnel.
- Looking forward meeting you around!



До зустрічі !

6. Outlook - FCC- hh and Flavours.

- The bb cross-section receives about a factor 5 enhancement at 100 TeV w.r.t. 14 TeV.
- The distinctive feature of FCC- hh is however that high- p_T Physics is enhanced by a far larger factor (~ 100).



Back


- It was still an early stage to devise a Flavour Physics case for the FCC- hh in the CDR. It will be part of the next stage of the Study.
- The progresses in data acquisition and triggering systems of the LHCb upgrades (to cope with high pile-up) will be invaluable in that respect.

7) References:

- CDR(s):
 - <https://fcc-cdr.web.cern.ch>
- FAQs about FCC:
 - <https://arxiv.org/pdf/1906.02693.pdf>

00) Back-up: Vertexing

- One of the most demanding requirement for vertex detectors comes from the missing momentum reconstruction inferred from the decay flight distances.
- Example: $X \rightarrow Y (Y \rightarrow [a]b) Z$ with a not reconstructed.



Back

- Three momentum components to be searched for:
 - The measurement of X momentum direction fixes 2 d.o.f.
 - An additional constraint closes the system: m_Y or a tertiary vertex.
 - Usually, quadratic form of the constraints: solution up to an ambiguity.

4. Distinctive features of FCC-ee at Z pole (Flavours)





Attribute	$\Upsilon(4S)$	pp	Z^0
All hadron species		✓	✓
High boost		✓	✓
Enormous production cross-section		✓	
Negligible trigger losses	✓		✓
Low backgrounds	✓		✓
Initial energy constraint	✓		(✓)

Some first mitigation comments:

vs LHC the modest cross-section partly compensated by the exquisite luminosity.

vs KEKB the initial energy constraint is diluted by the quark fragmentation; incoherent production of the b quarks.

4) Heavy Flavours: some examples of studies.



$B^0 \rightarrow K^* e^- e^+$

LHCb $\sigma = 75 \text{ MeV}/c^2$

Belle $\sigma = 5 \text{ MeV}/c^2$
[arXiv: 1904.02440]



© D. Hill

© R. Aleksan

Measurement of CP violation with $B_s \rightarrow D_s K$

$$\int L dt = 150 \text{ ab}^{-1}$$

PDG: $\gamma = (71.1^{+4.6}_{-5.3})^\circ$



Result 3: $\delta(\rho) \approx 3.2 \times 10^{-3} (\text{stat.})$
 $\delta(\sin^2 \phi_{CKM}) \approx \delta(\sin^2 \gamma) \approx 5 \times 10^{-3} (\text{stat.}) \cong \delta(\gamma) \approx 0.4^\circ (\text{stat.})$

Potential statistical gain of factor 4-5 with $D_s^\pm \rightarrow K^* K^\pm, \phi \rho^\pm, \dots$ but background needs to be studied (see later)
 Additional potential gain (another factor ~ 2) with $B_c \rightarrow D_s^\pm K^\mp, D_s^\pm K^{*\mp}, D_s^\pm K^{*+}$, most modes including $\gamma(\text{s})$

Some others around here:
<https://indico.in2p3.fr/event/23012/>